

# AUTHOR INDEX

- Ackermann RT, 41  
 Adams N, 51  
 Ahn C, 115  
 Albers GW, 18  
 Aldous MB, 68  
 ALLHAT Officers and Coordinators  
 for the ALLHAT Collaborative  
 Research Group, 86  
 Alper BS, 56  
 American-Canadian Thrombosis Study  
 Group, 6  
 Amir L, 35  
 Ammirati F, 121  
 Anand SS, 8  
 Antonson DL, 24  
 Armstrong PW, 4, 70  
 Ashton L, 10  
 Asplund K, 45  
 Astin JA, 107  
 Atkins D, 106  
 Atkinson RP, 91  
 ATLANTIS Study Investigators, 18  
 ATLAS Study Group, 4  
 ATTRACT Study Group, 16  
 Awasthi S, 24
- Badgett R, 27, 105  
 Baime MJ, 107  
 Bantje Th A, 101  
 Barnett HJ, 71  
 Barry MJ, 29  
 Basinski A, 67  
 Bass EB, 60  
 Bastian L, 34  
 Beach SR, 37  
 Beale RJ, 9  
 Becker DM, 32  
 Beghe C, 85  
 Beilby J, 103  
 Belknap SM, 44  
 Benesch CG, 19  
 Beyth RJ, 8, 117  
 Bihari DJ, 9  
 Bilous RW, 111  
 Black DM, 14  
 Bloch R, 21  
 Boissel JP, 88  
 Borgaonkar MR, 95  
 Boyle M, 110  
 Bradley C, 97  
 Brand R, 119  
 Brant RF, 6  
 Breedveld F, 16  
 Brennan PJ, 96  
 Bronsky EA, 3  
 Bryg DJ, 9  
 Bull SB, 42  
 Burr ML, 52  
 Burschtin O, 1
- Calam J, 11  
 Califf RM, 120
- Campbell IA, 52  
 Canadian Trial of Atrial Fibrillation  
 Investigators, 88  
 Canadian Trial of Physiologic Pacing  
 Investigators, 89  
 Capes SE, 116  
 CAPITAL Study Investigators, 72  
 Carbajal R, 25  
 CARE Group, A-11-12 (No. 2)  
 Chaisson RE, 94  
 Chambers BR, 48  
 Charlesworth G, 37  
 Chastre J, 93  
 Chauvet X, 25  
 Cheah G, 7  
 Cheng EY, 50  
 Chessman A, 72  
 Chippendale T, 91  
 CIDS Investigators, 43  
 Clark WM, 18  
 Colford JM Jr, 97  
 Colivicchi F, 121  
 Connolly SJ, 43, 89  
 Cornelissen PJ, 101  
 Couderc S, 25  
 Crebolder HF, 59  
 Crilly M, A-15 (No. 2)  
 Cripps AW, 2  
 Croft P, 26  
 Cunningham C, 110
- D'Amico D, 105  
 Damoiseaux RA, 62  
 Dans AL, A-11-12 (No. 1)  
 Dans LF, A-11-12 (No. 1)  
 Datta M, 23  
 Dawes M, A-15 (No. 1)  
 de Bie RA, 27  
 de Melker RA, 62  
 de Vet HC, 27  
 De Weerd W, 108  
 de Witte LP, 59  
 Dekker JH, 112  
 Del Mar CB, 62  
 Delaney B, 63  
 Delucchi KL, 109  
 Denman AM, 16  
 Deyle GD, 57  
 Dibley MJ, 23  
 Diederiks JP, 59  
 Dixon AS, 26  
 Dolovich LR, 7  
 Donnan GA, 48  
 Dorian P, 88  
 Douketis JD, 7  
 Dubé C, 63  
 Dudley RA, 119  
 Duggan AE, 36  
 Dunbabin DW, 45  
 Dutch Renal Artery Stenosis Intervention  
 Cooperative Study Group, 92  
 Dutch Tiotropium Study Group, 101
- Eakin EG, 104  
 Edelman JM, 3  
 Eisenstein EL, 120  
 Eland ME, 101  
 Eliasziw M, 71  
 ELITE II Investigators, 87  
 Elliott C, 36  
 Engels EA, 31  
 Entwistle VA, 28  
 Eriksson H, 117  
 Ernst E, 13, 107  
 Estrada CA, 113  
 Exercise Study Group, 3
- Fagerberg B, 46  
 Fagon J-Y, 93  
 Fairbairn I, 100  
 Fardy JM, 95  
 Farkouh ME, 70  
 Farrell S, 66  
 Feagan B, 95  
 Felson DT, 58  
 FitzGerald JM, 51  
 Fleming MF, 75  
 Fonseca V, 111  
 Foxwell AR, 2  
 Fracture Intervention  
 Trial Research Group, 14  
 Franse LV, 112  
 Frediani F, 105  
 French MT, 75  
 Furlan A, 19
- Garcia-Aranda JA, 64  
 Gasowski J, 41  
 Geerts WH, 66  
 Genant HK, 15  
 Gent M, 43, 89  
 Gerstein HC, 116  
 Gibson N, 60  
 Ginsberg JS, 7  
 Glasgow RE, 104  
 Glasziou P, A-11-12 (No. 3)  
 Goel M, 65  
 Goldstein S, 46  
 Gomis R, 49  
 Gordin F, 94  
 Götzsche PC, 67  
 Grady D, 106  
 Griffin S, 112  
 Griffiths TL, 52  
 Guiscafré H, 68  
 Gulin JR, 58  
 Gunton RW, 71  
 Gutierrez PR, 77  
 Guyatt GH, 85
- Haddon J, 51  
 Hagspiel KD, 32  
 Halperin JL, 90  
 Hankey GJ, 45, 71

- Hansen JF, 114  
Hansson PO, 117  
Harkness E, 107  
**Harper D, 113**  
Harrington RA, 76  
Harris ST, 15  
Hart RG, 90  
**Harvey RL, 90**  
**Havranek EP, 87**  
Hay EM, 26  
Hayes T, 66  
**Hay-Smith J, 108**  
Heart and Estrogen/progestin  
Replacement Study Research Group, 106  
Heart Outcomes Prevention Evaluation  
Study Investigators, 47  
Heine RJ, 112  
**Helewa A, 57**  
Henderson NE, 57  
Herrington D, 106  
Hess EV, 22  
**Heyland DK, 9**  
Higashida R, 19  
Hilden J, 114  
Hiller KO, 12  
Hjalmarson A, 46  
**Hla A, 114**  
**Hlatky MA, 120**  
Hoes AW, 62  
**Hoffman RM, 47**  
Holbrook AM, 7  
**Holloway RG, 19**  
**Honig PK, 3**  
Horn J, 20  
Hosie G, 26  
**Hui G, A-16 (No. 3)**  
Hulka BS, 118  
Hull RD, 6  
**Hunt D, 17, 116**  
  
Jackson JL, 11, 53  
Jadad AR, 110  
Janowsky EC, 118  
Jarvis G, 30  
Johansen KL, 119  
John G, 99  
**Johnell O, 14**  
**Jones A, 52**  
Jones NP, 49  
Jones PW, 102  
  
Kataria S, 69  
Katz J, 60  
**Kaul P, 70**  
**Keenan SP, 50, 93**  
Kerr CR, 89  
Kim M, 110  
**King PD, 55**  
**King S, 99**  
Kinmonth AL, 74  
Kitchener HC, 30  
Kohnen R, 12  
  
Korducki L, 101  
Krijnen P, 92  
Kupper LL, 118  
Kvarnström J, 103  
  
**Lacasse Y, 102**  
Lan AJ, 97  
**Lancaster T, 56**  
**Langhorne P, 20**  
Laroche C, 100  
Larsen CT, 114  
Lau CY, 72  
Lau J, 31  
LaValley MP, 58  
**Law F, 109**  
**Lawrie SM, 13**  
Leone M, 105  
Levi D, 55  
Lewis M, 26  
Lewis PR, 56  
Lilford R, 30  
Limburg M, 20  
Lincoff AM, 76  
Liu H, 77  
Liu Z, 23  
**Locke KA, A-15 (No. 3)**  
Logan RF, 36  
Lonn E, 4  
**Luxenberg JS, 77**  
  
**Macdonald JB, 100**  
MacIntosh DG, 95  
Maini R, 16  
Malmberg K, 116  
Manola J, 29  
**Margo G, 110**  
Mark DB, 76  
**Marrie TJ, 2, 72**  
Martínez H, 68  
Martin-Hirsch P, 30  
Mason J, 17  
Masson C, 109  
**Masters PA, 59**  
Matchar DB, 34  
Matekel RL, 57  
Mathur SK, 65  
Matts JP, 94  
Mbori-Ngacha D, 99  
McAlindon TE, 58  
**McAlister FA, A-11-12 (No. 2)**  
McBride R, 90  
McCorry DC, 34  
McGinn T, 121  
McIntosh A, 17  
**McMillan DD, 25**  
**Mehra MR, 46**  
**Meisner JS, 114**  
**Melnikow J, 30, 34**  
MERIT-HF Study Group, 46  
Mermel LA, 96  
Messing EM, 29  
**Millard PS, 94**  
  
Miller RG, 54  
Milligan P, 98  
**Mills CS, 119**  
Milstein A, 119  
Mimouni M, 35  
**Minde K, 61**  
Mitchell JD, 54  
Moher D, 63  
Molloy DW, 85  
**Moody KA, 94**  
Moore DH, 54  
Morales-Ramirez J, 22  
Morton S, 21  
Moss H, 100  
Moyer VA, 115  
Mullins MD, 32  
**Mulrow CD, 10, 41**  
Mundt MP, 75  
Muñoz AE, 55  
Muñoz O, 68  
  
Nduati R, 99  
Nelemans PJ, 27  
Nevitt MC, 14  
Newby LK, 120  
Nielsen OW, 114  
Niemelä M, 44  
**Novak F, 9**  
**Nuovo J, 30, 34**  
Nurko S, 64  
  
OESIL Study Investigators, 121  
Ohl CA, 113  
O'Keeffe C, 17  
Olivier-Martin M, 25  
Olsen O, 67  
O'Malley PG, 11, 53  
Ornstein MP, 42  
Ostermann ME, 50  
Otten N, 63  
  
Packer M, 4  
**Pahor M, 86**  
Palafox M, 68  
Paramothayan NS, 102  
Patel SC, 113  
Patel SR, 113  
Paterson SM, 26  
Patwardhan R, 49, 111  
**Pelosi AJ, 10**  
**Perlman M, 115**  
**Peters J, 1**  
**Peterson WL, 36**  
Peuhkurinen K, 44  
Philbrick JT, 32  
Philipp M, 12  
Phillips MR, 23  
Pieterman H, 92  
Pinder M, 98  
Pitt B, 87  
Pittler MH, 13  
Podestá A, 55

- Polashenski WA, A-11-12 (No. 3)  
 Poole-Wilson PA, 4, 87  
 Poussa T, 103  
 Price J, 53  
 Price RN, 98  
 PROACT Investigators, 19  
 Psaty BM, 86  
 Pugh JA, 49  
 Pulicino P, 91  
 Puntis JW, 64
- Raco D, 76  
 Rajesh VT, 69  
 Ramchandani P, 61  
 Raskob GE, 6, 33  
 Rathbun SW, 33  
 Ray J, 42  
 Reginster J-Y, 58  
 Reiner A, 5  
 Rennie DJ, 119  
 Reyes H, 68  
 Richardson WS, A-11-12 (No. 3)  
 Riley KM, 104  
 Roberts RS, 43  
 Rodrigo C, 1  
 Rodrigo G, 1  
 Roger VL, 70  
 Rosenberg W, A-16 (No. 1)  
 Rosenstock J, 111  
 Roy D, 88  
 Russo R, 85  
 Rutter C, 73
- Sackett DL, A-11-12 (No. 2)  
 Salzman A, 111  
 Santini M, 121  
 Santoro J, 53  
 Sarkisian CA, 77  
 Sarosdy M, 29  
 Schachar R, 110  
 Schein OD, 60  
 Schmitt BP, 6  
 Schuger CD, 43  
 Schulz R, 37  
 Scott JT, 28  
 Sees KL, 109  
 Segal R, 87  
 Seiferling RA, 50  
 Selby R, 66  
 Shah SR, 65  
 Shaw M, 66  
 Shekelle P, 21  
 Sherman DG, 91
- Sherman SE, 104  
 Shikare S, 65  
 Shragge D, 51  
 Shridharani K, 121  
 Shukla VK, 63  
 Sibbald WJ, 50  
 Silverman WB, 65  
 Silverton SF, 15  
 Simon GE, 73  
 Singhi S, 69  
 Smith DA, 116  
 Smith RC, 28  
 Sneed S, 115  
 Soonawalla ZF, 65  
 Sörbo J, 117  
 Sowden AJ, 28  
 Squatrito S, 49  
 St. Clair EW, 16  
 Staessen JA, 41  
 Stanbrook MB, 33  
 Staszewski S, 22  
 STAT Participants, 91  
 Stenius-Aarniala B, 103  
 Stevens L, 74  
 Stores G, 61  
 Straus SE, A-11-12 (No. 2)  
 Strong MJ, 54  
 Study 006 Team, 22  
 Study of Medical Testing for  
 Cataract Surgery, 60  
 Study of Osteoporotic Fractures  
 Research Group, 77  
 Sturmans F, 27  
 Sudbury P, 75  
 Sudlow CL, 45  
 Svanegaard J, 114  
 Swainson CR, 92  
 Symmons DPM, 118  
 SYMPHONY Investigators, 5
- Talajic M, 88  
 Tashima KT, 22  
 Taylor PR, 48  
 Thompson C, 74  
 Thompson DE, 14  
 Tomkins G, 11  
 Trivedi M, 10  
 Turpin JA, 3  
 Tyrer P, 74
- Valk GD, 112  
 van Balen FA, 62  
 van Eijk JT, 112
- van Haastregt JC, 59  
 van Jaarsveld BC, 92  
 Van Kampen M, 108  
 van Noord JA, 101  
 Van Poppel H, 108  
 van Rossum E, 59  
 Vanderhoof JA, 24  
 VAP Trial Group, 93  
 Verheij TJ, 62  
 Vermeire E, 12  
 Vertebral Efficacy with Risedronate  
 Therapy (VERT) Study Group, 15  
 Vickrey BG, 21  
 von Dadelzen P, 42  
 von Seidlein L, 98  
 VonKorff M, 73
- Wagner E, 73  
 Waisman Y, 35  
 Wan C, 23  
 Wang JG, 41  
 Watt I, 28  
 Watts NB, 15  
 Webb V, 61  
 Wechsler L, 19  
 Weiss R, 89  
 Wenger NK, 106  
 Weston SA, 70  
 Wheeler SL, 72  
 Whitney DB, 24  
 Whitsett TL, 33  
 Wiedemann HP, 101  
 Wiese W, 113  
 Wiggs L, 61  
 Wigton RS, 35  
 Willett LR, 31  
 Williams JW Jr, 10, 73  
 Wilson MC, A-11-12 (No. 3)  
 Wissman S, 18  
 Wolff M, 93  
 Wolffenbuttel BHR, 49  
 Worona LB, 64
- Ylitalo A, 44  
 You RX, 48  
 Yusuf S, 8
- Zerem E, 35  
 Zlochisty O, 64  
 Zucker D, 31

## A

- Abdominal pain
  - Review: Antidepressive agents improve symptoms and reduce pain in patients with functional gastrointestinal disorders, 11
- Accident prevention
  - Review: Evidence is unclear that preventive home visits for elderly persons in the community improve mortality or health outcomes, 59
- Acetates
  - Oral montelukast was better than inhaled salmeterol for reducing exercise-induced bronchoconstriction in adults with asthma, 3
- Activities of daily living
  - A clinical prediction tool that includes modifiable risk factors predicted functional decline in elderly women, 77
- Acyclovir
  - Review: Early treatment of acute herpes zoster may prevent or shorten the duration of postherpetic neuralgia, 56
- Administration, oral
  - Oral amiodarone was effective and safe for recent-onset atrial fibrillation, 44
- Adrenal cortex hormones
  - Review: Oral but not inhaled corticosteroids improve radiographic outcomes in patients with pulmonary sarcoidosis, 102
- Adrenergic  $\beta$ -agonists
  - Review: Ipratropium bromide with  $\beta$ -agonists improves pulmonary function and reduces hospitalizations in acute asthma, 1
- Adrenergic  $\beta$ -antagonists
  - Controlled- and extended-release metoprolol reduced death, hospital admissions, and symptoms in chronic heart failure, 46
- Advance directives
  - An advance directive program in nursing homes reduced health services utilization without affecting patient satisfaction, 85
- Aged
  - A clinical prediction tool that includes modifiable risk factors predicted functional decline in elderly women, 77
  - Losartan did not improve survival more than did captopril but was better tolerated in symptomatic heart failure, 87
  - Review: Evidence is unclear that preventive home visits for elderly persons in the community improve mortality or health outcomes, 59
  - Review: Isolated systolic hypertension increases mortality and morbidity in elderly persons and should be treated, 41
- AIDS-related opportunistic infections
  - Rifampin and pyrazinamide for 2 months prevented TB in patients with HIV infection who had a positive tuberculin test result, 94
- Albuterol
  - Oral montelukast was better than inhaled salmeterol for reducing exercise-induced bronchoconstriction in adults with asthma, 3
- Alcoholism
  - Brief physician advice to problem drinkers reduced alcohol intake and societal costs, 75
- Alendronate
  - Alendronate reduced days of bed rest and limited activity in postmenopausal women with osteoporosis and existing fractures, 14
- Algorithms
  - A new arrival: evidence about differential diagnosis (Editorial), A-11-12 (No. 3)
  - A simple algorithm improved diagnostic performance for patients presenting with syncope, 121
- Alternative medicine
  - Review: "Distant healing" is often effective for improving patient outcomes, 107
- Ambulatory care
  - An outpatient multidisciplinary pulmonary rehabilitation program was effective in disabling chronic lung disease, 52
- Amiodarone
  - The implantable cardioverter defibrillator did not reduce mortality in ventricular arrhythmia, 43
  - Low-dose amiodarone was better than sotalol or propafenone for preventing first recurrence of atrial fibrillation, 88
  - Oral amiodarone was effective and safe for recent-onset atrial fibrillation, 44
- Amoxicillin
  - Amoxicillin resolved otitis media symptoms in young children better than did placebo at day 4 but not at day 11, 62
  - Review: Twice-daily dosing of penicillin V is as effective as more frequent dosing for streptococcal tonsillopharyngitis, 97
- Amyotrophic lateral sclerosis
  - Review: Riluzole increases 12-month survival in amyotrophic lateral sclerosis, 54
- Analgesics
  - Pacifiers have greater analgesic effect than sweet solutions during venipuncture in full-term newborns, 25
- Ancrod
  - More patients with acute ischemic stroke who received ancrod had good functional status at 3 months, 91
- Angina, unstable
  - Eptifibatide was cost-effective in patients with unstable angina or non-Q-wave myocardial infarction, 76
  - Sibrafiban was more toxic than aspirin for prevention of cardiovascular events after acute coronary syndromes, 5
  - Women with unstable angina received fewer cardiac procedures than did men but were less likely to develop other cardiac events, 70
- Angioplasty, balloon
  - Angioplasty was no better than medication for reducing blood pressure or slowing disease progression in renal artery stenosis, 92
- Angiotensin II
  - Losartan did not improve survival more than did captopril but was better tolerated in symptomatic heart failure, 87
- Angiotensin-converting enzyme inhibitors
  - High-dose lisinopril was more effective than low-dose for reducing combined mortality and cardiovascular events in CHF, 4
- Anoxia
  - A respiratory rate of  $\geq 60$  breaths per minute had high sensitivity for detecting hypoxia in infants, 69
- Anti-arrhythmia agents
  - Oral amiodarone was effective and safe for recent-onset atrial fibrillation, 44
- Antibiotics
  - Lactobacillus GG* reduced diarrhea incidence in children treated with antibiotics, 24
  - Review: Antimicrobial prophylaxis reduces asymptomatic bacteriuria in patients with neurogenic bladder, 21
- Anticoagulants
  - The recurrence rate of venous thromboembolism after a first or second episode of deep venous thrombosis was high, 117
  - Review: High- and moderate-dose oral anticoagulants reduce events in CAD but increase major bleeding and are no more effective than aspirin, 8

## Antidepressants

- Review: Antidepressive agents improve symptoms and reduce pain in patients with functional gastrointestinal disorders, 11
- Review: Antidepressants are effective for clinical improvement in unexplained physical symptoms and syndromes, 53
- Review: Newer and older antidepressants have similar efficacy and total discontinuation rates but different side effects, 10

## Antihypertensive agents

- Angioplasty was no better than medication for reducing blood pressure or slowing disease progression in renal artery stenosis, 92
- Doxazosin was associated with more stroke and cardiovascular disease events than chlorthalidone in high-risk hypertension, 86
- Review: Isolated systolic hypertension increases mortality and morbidity in elderly persons and should be treated, 41
- Review: Treatment-induced blood pressure reductions in pregnancy may be associated with decreased fetal growth, 42

## Anti-inflammatory agents, nonsteroidal

- Ketoprofen given twice daily and interferon- $\alpha$  2b increased and maintained response in chronic HCV-related liver disease, 55

## Antitubercular agents

- Review: Antimycobacterial therapy does not increase the rate of maintenance of remission in Crohn disease, 95
- Rifampin and pyrazinamide for 2 months prevented TB in patients with HIV infection who had a positive tuberculin test result, 94

## Anxiety disorders

- Review: Kava extract reduces symptoms in adults with anxiety, 13

## Artesunate

- Adding 3-dose artesunate to pyrimethamine-sulfadoxine reduced treatment failure in children with acute uncomplicated malaria, 98

## Arthritis, rheumatoid

- Infliximab at different doses was effective for rheumatoid arthritis after unsuccessful methotrexate treatment, 16

## Aspirin

- Review: Aspirin lowers the risk for MI but not for stroke in persons who do not have established vascular disease, 90
- Review: High- and moderate-dose oral anticoagulants reduce events in CAD but increase major bleeding and are no more effective than aspirin, 8
- Review: Thienopyridines are more effective than aspirin for preventing vascular events in high-risk patients, 45
- Sibrafiban was more toxic than aspirin for prevention of cardiovascular events after acute coronary syndromes, 5

## Asthma

- Inhaled budesonide and prednisone led to similar rates of asthma relapse after emergency department discharge, 51
- Oral montelukast was better than inhaled salmeterol for reducing exercise-induced bronchoconstriction in adults with asthma, 3
- Review: Ipratropium bromide with  $\beta$ -agonists improves pulmonary function and reduces hospitalizations in acute asthma, 1
- Weight loss in obese patients with asthma improved lung function and health status, 103

## Atherosclerosis

- Angioplasty was no better than medication for reducing blood pressure or slowing disease progression in renal artery stenosis, 92
- Review: Aspirin lowers the risk for MI but not for stroke in persons who do not have established vascular disease, 90

## Atrial fibrillation

- Low-dose amiodarone was better than sotalol or propafenone for preventing first recurrence of atrial fibrillation, 88

Oral amiodarone was effective and safe for recent-onset atrial fibrillation, 44

Physiologic and ventricular pacing had similar rates of combined stroke and cardiovascular mortality, 89

## Attention deficit disorder with hyperactivity

- Review: Pharmacologic interventions are more effective than nonpharmacologic for attention-deficit hyperactivity disorder, 110

## Autoimmune diseases

- Review: Silicone breast implants do not increase the risk for connective-tissue diseases, 118

## B

## Back pain

- Alendronate reduced days of bed rest and limited activity in postmenopausal women with osteoporosis and existing fractures, 14

## Bacteremia

- Review: Intravascular catheter-related infections can be prevented by using simple procedures, 96

## Bed rest

- Alendronate reduced days of bed rest and limited activity in postmenopausal women with osteoporosis and existing fractures, 14

## Behavior therapy

- Review: Long-term effectiveness of nondrug treatments for sleep problems in young children is more evident, 61

## Bilirubin

- Clinical examination could not accurately predict neonatal jaundice, 115

## Biofeedback

- Pelvic floor re-education reduced incontinence at 1 year after radical prostatectomy, 108

## Birth weight

- Review: Treatment-induced blood pressure reductions in pregnancy may be associated with decreased fetal growth, 42

## Bladder, neurogenic

- Review: Antimicrobial prophylaxis reduces asymptomatic bacteriuria in patients with neurogenic bladder, 21

## Blood glucose

- Low-dose rosiglitazone plus sulfonylurea was effective and safe for treating hyperglycemia in type 2 diabetes mellitus, 49

## Bone density

- Risedronate was effective and well tolerated in postmenopausal women with osteoporosis, 15

## Bottle feeding

- Risk for mother-to-child HIV-1 infection was increased by breast-feeding, 99

## Bradycardia

- Physiologic and ventricular pacing had similar rates of combined stroke and cardiovascular mortality, 89

## Brain ischemia

- Review: Calcium antagonists do not reduce death or dependency in acute ischemic stroke, 20

## Breast feeding

- Risk for mother-to-child HIV-1 infection was increased by breast-feeding, 99

## Breast implants

- Review: Silicone breast implants do not increase the risk for connective-tissue diseases, 118

## Breast neoplasms

- Review: High-quality studies do not show reductions in breast cancer mortality for women screened with mammography, 67

Bronchitis

Review: Oral *Haemophilus influenzae* vaccination reduces the number and severity of recurrent bronchitis episodes, 2

Bronchoconstriction

Oral montelukast was better than inhaled salmeterol for reducing exercise-induced bronchoconstriction in adults with asthma, 3

Bronchoscopy

CT scanning before fiberoptic bronchoscopy improved the accuracy of bronchoscopy in suspected endobronchial carcinoma, 100

Budesonide

Inhaled budesonide and prednisone led to similar rates of asthma relapse after emergency department discharge, 51

C

Calcium channel blockers

Review: Calcium antagonists do not reduce death or dependency in acute ischemic stroke, 20

Captopril

Losartan did not improve survival more than did captopril but was better tolerated in symptomatic heart failure, 87

Cardiac pacing, artificial

Physiologic and ventricular pacing had similar rates of combined stroke and cardiovascular mortality, 89

Caregivers

Older adults who reported strain when caring for a spouse with disabilities had increased mortality, 37

Carotid stenosis

Patients with carotid artery stenosis had a high risk for stroke, and some types of stroke were not prevented by endarterectomy, 71  
Review: Carotid endarterectomy for asymptomatic carotid stenosis does not reduce stroke or death, 48

Cataract

Routine testing before cataract surgery did not reduce medical adverse events, 60

Catheterization, central venous

Review: Intravascular catheter-related infections can be prevented by using simple procedures, 96

Catheters, indwelling

Review: Intravascular catheter-related infections can be prevented by using simple procedures, 96

Cerebrovascular accident

Intra-arterial recombinant prourokinase improved neurologic recovery in acute middle cerebral artery ischemic stroke, 19  
More patients with acute ischemic stroke who received anecro had good functional status at 3 months, 91  
Patients with carotid artery stenosis had a high risk for stroke, and some types of stroke were not prevented by endarterectomy, 71  
Physiologic and ventricular pacing had similar rates of combined stroke and cardiovascular mortality, 89  
Recombinant tissue-type plasminogen activator at 3 to 5 hours after ischemic acute stroke onset was not effective or safe, 18  
Review: Aspirin lowers the risk for MI but not for stroke in persons who do not have established vascular disease, 90  
Review: Thienopyridines are more effective than aspirin for preventing vascular events in high-risk patients, 45

Cervical intraepithelial neoplasia

Review: The extended-tip spatula is more effective than the Ayre's spatula for detecting karyosis in cervical smears, 30  
Review: Newer technologies improve sensitivity in detecting cervical lesions but at a substantially increased cost, 34

Child

Review: Long-term effectiveness of nondrug treatments for sleep problems in young children is more evident, 61  
Tachypnea is a useful predictor of pneumonia in children with acute respiratory infection, 68

Chlorthalidone

Doxazosin was associated with more stroke and cardiovascular disease events than chlorthalidone in high-risk hypertension, 86

Cholangiography

Biliary scintiscan had high sensitivity and specificity for predicting pathologic findings in the common bile duct, 65

Cholelithiasis

Biliary scintiscan had high sensitivity and specificity for predicting pathologic findings in the common bile duct, 65

Chondroitin

Review: Glucosamine and chondroitin improve outcomes in osteoarthritis, but the magnitude of effect is unclear, 58

Chronic disease

Review: Antidepressants are effective for clinical improvement in unexplained physical symptoms and syndromes, 53

Cisapride

Cisapride was effective in the short term for constipation in children, 64

Review: Cisapride reduces overall and global symptoms in adults with nonulcer dyspepsia, but study quality is poor, 63

Cluster headache

Verapamil reduced attack frequency and the use of abortive agents in episodic cluster headache, 105

Colonic diseases, functional

Review: Antidepressive agents improve symptoms and reduce pain in patients with functional gastrointestinal disorders, 11

Community-acquired infections

A critical pathway reduced resource use without compromising safety and effectiveness in community-acquired pneumonia, 72

Constipation

Cisapride was effective in the short term for constipation in children, 64

Counseling

Brief physician advice to problem drinkers reduced alcohol intake and societal costs, 75

Critical illness

Review: Enteral immunonutrition reduces infection risk, days on ventilation, and hospital stay in critically ill patients, 9

Critical pathways

A critical pathway reduced resource use without compromising safety and effectiveness in community-acquired pneumonia, 72

Crohn disease

Review: Antimycobacterial therapy does not increase the rate of maintenance of remission in Crohn disease, 95

Cytology

Review: Newer technologies improve sensitivity in detecting cervical lesions but at a substantially increased cost, 34

D

Databases, bibliographic

PDQ Evidence-Based Principles and Practice (Resource Corner), A-16 (No. 3)

Defibrillators, implantable

The implantable cardioverter defibrillator did not reduce mortality in ventricular arrhythmia, 43



## Depression

- Feedback to physicians plus telephone care management improved outcomes in primary care patients with depression, 73
- Hypericum* extract was better than placebo and equivalent to imipramine for moderate depression, 12
- Practice-based education did not increase recognition of depression by primary-care physicians nor improve the outcome of depression, 74
- Review: Newer and older antidepressants have similar efficacy and total discontinuation rates but different side effects, 10

## Developing countries

- The need and means for evidence-based medicine in developing countries (Editorial), A-11-12 (No. 1)

## Dextroamphetamine

- Review: Pharmacologic interventions are more effective than non-pharmacologic for attention-deficit hyperactivity disorder, 110

## Diabetes mellitus

- Low-dose rosiglitazone plus sulfonylurea was effective and safe for treating hyperglycemia in type 2 diabetes mellitus, 49
- Metformin plus rosiglitazone had more effective glycemic control than metformin alone in type 2 diabetes mellitus, 111
- Neuropathic sensory symptoms did not accurately detect polyneuropathy in type 2 diabetes mellitus, 112
- Review: Hyperglycemia after myocardial infarction increases the risk for death in patients with and without diabetes mellitus, 116
- Review: Inadequate data exist on prevention and treatment strategies for foot ulcer in diabetes mellitus, 17

## Diagnosis, differential

- A new arrival: evidence about differential diagnosis (Editorial), A-11-12 (No. 3)
- A simple algorithm improved physicians' diagnostic performance for patients presenting with syncope, 121

## Diagnostic tests, routine

- Routine testing before cataract surgery did not reduce medical adverse events, 60

## Diarrhea

- Lactobacillus GG* reduced diarrhea incidence in children treated with antibiotics, 24
- Low-volume, high-frequency cow's milk feedings sped recovery from diarrhea in infants, 23

## Disease transmission, vertical

- Risk for mother-to-child HIV-1 infection was increased by breast-feeding, 99

## Doxazosin

- Doxazosin was associated with more stroke and cardiovascular disease events than chlorthalidone in high-risk hypertension, 86

## Drug therapy

- Review: Long-term effectiveness of nondrug treatments for sleep problems in young children is more evident, 61

## Dyspepsia

- Review: Antidepressive agents improve symptoms and reduce pain in patients with functional gastrointestinal disorders, 11
- Review: Cisapride reduces overall and global symptoms in adults with nonulcer dyspepsia, but study quality is poor, 63

## E

## Echocardiography/electrocardiography

- A combination of tests done in general practice could assess the risk for left ventricular systolic dysfunction, 114

## Endarterectomy, carotid

- Review: Carotid endarterectomy for asymptomatic carotid stenosis does not reduce stroke or death, 48

## Enteral nutrition

- Review: Enteral immunonutrition reduces infection risk, days on ventilation, and hospital stay in critically ill patients, 9

## Estrogen replacement therapy

- Hormone replacement therapy was associated with increased venous thromboembolism and deep venous thrombosis, 106

## Etidronate disodium, analogs and derivatives

- Risedronate was effective and well tolerated in postmenopausal women with osteoporosis, 15

## Evidence-based medicine

- Bandolier (Resource Corner), A-15 (No. 2)
- Best Evidence 4 (Resource Corner), A-15 (No. 3)
- The need and means for evidence-based medicine in developing countries (Editorial), A-11-12 (No. 1)
- PDQ Evidence-Based Principles and Practice* (Resource Corner), A-16 (No. 3)

## Exercise

- Manual physical therapy and exercise improved function in osteoarthritis of the knee, 57
- Review: Brief primary care interventions are moderately effective for increasing physical activity, 104

## F

## Feedback

- Feedback to physicians plus telephone care management improved outcomes in primary care patients with depression, 73

## Fibrinogen degradation products

- SimpliRED D-dimer assay did not rule out venous thromboembolism in the emergency department, 66

## Fibrinolytic agents

- More patients with acute ischemic stroke who received anistreplase had good functional status at 3 months, 91

## Fractures

- Risedronate was effective and well tolerated in postmenopausal women with osteoporosis, 15

## G

## Glucosamine

- Review: Glucosamine and chondroitin improve outcomes in osteoarthritis, but the magnitude of effect is unclear, 58

## Goserelin

- Antiandrogen therapy immediately after surgery for prostate cancer improved survival, 29

## H

*Haemophilus* vaccines

- Review: Oral *Haemophilus influenzae* vaccination reduces the number and severity of recurrent bronchitis episodes, 2

## Health care costs

- An advance directive program in nursing homes reduced health services utilization without affecting patient satisfaction, 85

## Health services, utilization

- An outpatient multidisciplinary pulmonary rehabilitation program was effective in disabling chronic lung disease, 52

Heart failure, congestive  
 Controlled- and extended-release metoprolol reduced death, hospital admissions, and symptoms in chronic heart failure, 46  
 High-dose lisinopril was more effective than low-dose for reducing combined mortality and cardiovascular events in CHF, 4

*Helicobacter* infections  
 The near-patient test FlexSure had low sensitivity and high specificity for detecting *Helicobacter pylori* infection, 36

Heparin  
 Low-molecular-weight heparin reduced recurrent VTE in patients with pulmonary embolism and proximal DVT, 6  
 Review: Low-molecular-weight heparin reduces death but not recurrent VTE events, bleeding, or thrombocytopenia in VTE, 7

Hepatitis C, chronic  
 Ketoprofen given twice daily and interferon- $\alpha$  2b increased and maintained response in chronic HCV-related liver disease, 55

Herpes zoster  
 Review: Early treatment of acute herpes zoster may prevent or shorten the duration of postherpetic neuralgia, 56

HIV infections  
 Efavirenz plus zidovudine and lamivudine was more effective than indinavir plus zidovudine and lamivudine in adults with HIV-1 infection, 22  
 Rifampin and pyrazinamide for 2 months prevented TB in patients with HIV infection who had a positive tuberculin test result, 94  
 Risk for mother-to-child HIV-1 infection was increased by breast-feeding, 99

HIV protease inhibitors  
 Efavirenz plus zidovudine and lamivudine was more effective than indinavir plus zidovudine and lamivudine in adults with HIV-1 infection, 22

Home nursing  
 Older adults who reported strain when caring for a spouse with disabilities had increased mortality, 37

Hospital mortality  
 Review: High-volume hospitals had lower hospital mortality rates than did low-volume hospitals for various procedures, 119

Hospitalization  
 A critical pathway reduced resource use without compromising safety and effectiveness in community-acquired pneumonia, 72

House calls  
 Review: Evidence is unclear that preventive home visits for elderly persons in the community improve mortality or health outcomes, 59

Hyperglycemia  
 Review: Hyperglycemia after myocardial infarction increases the risk for death in patients with and without diabetes mellitus, 116

*Hypericum* extract  
*Hypericum* extract was better than placebo and equivalent to imipramine for moderate depression, 12

Hypertension  
 Angioplasty was no better than medication for reducing blood pressure or slowing disease progression in renal artery stenosis, 92  
 Review: Isolated systolic hypertension increases mortality and morbidity in elderly persons and should be treated, 41  
 Review: Treatment-induced blood pressure reductions in pregnancy may be associated with decreased fetal growth, 42

Hypnotics and sedatives  
 Review: Quality of sedation in the ICU is similar for propofol and midazolam, but propofol shortens the time to extubation, 50

Hypoglycemic agents  
 Low-dose rosiglitazone plus sulfonylurea was effective and safe for treating hyperglycemia in type 2 diabetes mellitus, 49  
 Metformin plus rosiglitazone had more effective glycemic control than metformin alone in type 2 diabetes mellitus, 111

## I

Imipramine  
*Hypericum* extract was better than placebo and equivalent to imipramine for moderate depression, 12

Immune system  
 Review: Enteral immunonutrition reduces infection risk, days on ventilation, and hospital stay in critically ill patients, 9

Indinavir  
 Efavirenz plus zidovudine and lamivudine was more effective than indinavir plus zidovudine and lamivudine in adults with HIV-1 infection, 22

Infant  
 A respiratory rate of  $\geq 60$  breaths per minute had high sensitivity for detecting hypoxia in infants, 69

Infection control  
 Review: Intravascular catheter-related infections can be prevented by using simple procedures, 96

Infliximab  
 Infliximab at different doses was effective for rheumatoid arthritis after unsuccessful methotrexate treatment, 16

Injections  
 Local injection, naproxen, and simple analgesia led to similar 1-year rates of symptom relief in lateral epicondylitis, 26  
 Review: Injection therapy is not better than placebo for relieving pain in benign chronic low-back pain, 27

Intensive care  
 Invasive diagnosis for ventilator-associated pneumonia reduced 14-day mortality and antibiotic use, 93  
 Review: Enteral immunonutrition reduces infection risk, days on ventilation, and hospital stay in critically ill patients, 9  
 Review: Quality of sedation in the ICU is similar for propofol and midazolam, but propofol shortens the time to extubation, 50

Interferon,  $\alpha$  2b  
 Ketoprofen given twice daily and interferon- $\alpha$  2b increased and maintained response in chronic HCV-related liver disease, 55

Intracranial hemorrhages  
 Review: Aspirin lowers the risk for MI but not for stroke in persons who do not have established vascular disease, 90

Ipratropium  
 Review: Ipratropium bromide with  $\beta$ -agonists improves pulmonary function and reduces hospitalizations in acute asthma, 1  
 Tiotropium improved lung function more than did ipratropium in chronic obstructive pulmonary disease, 101

Isoniazid  
 Rifampin and pyrazinamide for 2 months prevented TB in patients with HIV infection who had a positive tuberculin test result, 94

## J

Jaundice  
 Biliary scintiscan had high sensitivity and specificity for predicting pathologic findings in the common bile duct, 65  
 Clinical examination could not accurately predict neonatal jaundice, 115



## K

- Kava  
Review: Kava extract reduces symptoms in adults with anxiety, 13
- Ketoprofen  
Ketoprofen given twice daily and interferon- $\alpha$  2b increased and maintained response in chronic HCV-related liver disease, 55

## L

- Lactobacillus casei*  
*Lactobacillus GG* reduced diarrhea incidence in children treated with antibiotics, 24
- Leukotriene antagonists  
Oral montelukast was better than inhaled salmeterol for reducing exercise-induced bronchoconstriction in adults with asthma, 3
- Lisinopril  
High-dose lisinopril was more effective than low-dose for reducing combined mortality and cardiovascular events in CHF, 4
- Low-back pain  
Review: Injection therapy is not better than placebo for relieving pain in benign chronic low-back pain, 27
- Lung diseases, obstructive  
An outpatient multidisciplinary pulmonary rehabilitation program was effective in disabling chronic lung disease, 52  
Tiotropium improved lung function more than did ipratropium in chronic obstructive pulmonary disease, 101
- Lung neoplasms  
CT scanning before fiberoptic bronchoscopy improved the accuracy of bronchoscopy in suspected endobronchial carcinoma, 100

## M

- Malaria  
Adding 3-dose artesunate to pyrimethamine-sulfadoxine reduced treatment failure in children with acute uncomplicated malaria, 98
- Mammography  
Review: High-quality studies do not show reductions in breast cancer mortality for women screened with mammography, 67
- Maxillary sinusitis  
Review: Diagnostic tests for bacterial rhinosinusitis have moderate accuracy; antibiotics reduce clinical failures, 31
- Mental healing  
Review: "Distant healing" is often effective for improving patient outcomes, 107
- Metformin  
Metformin plus rosiglitazone had more effective glycemic control than metformin alone in type 2 diabetes mellitus, 111
- Methadone  
Methadone maintenance was more effective for treatment retention of opioid dependence than was psychosocially enriched detoxification, 109
- Methylphenidate  
Review: Pharmacologic interventions are more effective than non-pharmacologic for attention-deficit hyperactivity disorder, 110
- Methylprednisolone  
Local injection, naproxen, and simple analgesia led to similar 1-year rates of symptom relief in lateral epicondylitis, 26
- Metoprolol  
Controlled- and extended-release metoprolol reduced death, hospital admissions, and symptoms in chronic heart failure, 46

## Milk

- Low-volume, high-frequency cow's milk feedings sped recovery from diarrhea in infants, 23

## Mortality

- Older adults who reported strain when caring for a spouse with disabilities had increased mortality, 37

## Mycobacterium infections, atypical

- Review: Antimycobacterial therapy does not increase the rate of maintenance of remission in Crohn disease, 95

## Myocardial infarction

- Eptifibatide was cost-effective in patients with unstable angina or non-Q-wave myocardial infarction, 76  
Hospitalization for more than 3 days after thrombolysis for uncomplicated myocardial infarction was not cost-effective, 120  
Losartan did not improve survival more than did captopril but was better tolerated in symptomatic heart failure, 87  
Review: Hyperglycemia after myocardial infarction increases the risk for death in patients with and without diabetes mellitus, 116

## N

## Naproxen

- Local injection, naproxen, and simple analgesia led to similar 1-year rates of symptom relief in lateral epicondylitis, 26

## Narcotics

- Methadone maintenance was more effective for treatment retention of opioid dependence than was psychosocially enriched detoxification, 109

## Neoplasms

- Review: Providing recordings or summaries of consultations may help patients with cancer, 28

## Neuralgia

- Review: Early treatment of acute herpes zoster may prevent or shorten the duration of postherpetic neuralgia, 56

## Nursing homes

- An advance directive program in nursing homes reduced health services utilization without affecting patient satisfaction, 85

## O

## Obesity

- Weight loss in obese patients with asthma improved lung function and health status, 103

## Opioid-related disorders

- Methadone maintenance was more effective for treatment retention of opioid dependence than was psychosocially enriched detoxification, 109

## Orchiectomy

- Antiandrogen therapy immediately after surgery for prostate cancer improved survival, 29

## Osteoarthritis

- Manual physical therapy and exercise improved function in osteoarthritis of the knee, 57

- Review: Glucosamine and chondroitin improve outcomes in osteoarthritis, but the magnitude of effect is unclear, 58

## Osteoporosis, postmenopausal

- Alendronate reduced days of bed rest and limited activity in postmenopausal women with osteoporosis and existing fractures, 14

## Otitis media

- Amoxicillin resolved otitis media symptoms in young children better than did placebo at day 4 but not at day 11, 62

Oxazines

Efavirenz plus zidovudine and lamivudine was more effective than indinavir plus zidovudine and lamivudine in adults with HIV-1 infection, 22

Oximes

Sibrafiban was more toxic than aspirin for prevention of cardiovascular events after acute coronary syndromes, 5

P

Pacifiers

Pacifiers have greater analgesic effect than sweet solutions during venipuncture in full-term newborns, 25

Pain

Review: Antidepressants are effective for clinical improvement in unexplained physical symptoms and syndromes, 53

Review: Injection therapy is not better than placebo for relieving pain in benign chronic low-back pain, 27

Pancreatitis

Biliary scintiscan had high sensitivity and specificity for predicting pathologic findings in the common bile duct, 65

Patient discharge

Hospitalization for more than 3 days after thrombolysis for uncomplicated myocardial infarction was not cost-effective, 120

Patient education

An outpatient multidisciplinary pulmonary rehabilitation program was effective in disabling chronic lung disease, 52

Review: Brief primary care interventions are moderately effective for increasing physical activity, 104

Review: Inadequate data exist on prevention and treatment strategies for foot ulcer in diabetes mellitus, 17

Patient satisfaction

An advance directive program in nursing homes reduced health services utilization without affecting patient satisfaction, 85

Pelvic floor

Pelvic floor re-education reduced incontinence at 1 year after radical prostatectomy, 108

Pemoline

Review: Pharmacologic interventions are more effective than non-pharmacologic for attention-deficit hyperactivity disorder, 110

Penicillins

Review: Twice-daily dosing of penicillin V is as effective as more frequent dosing for streptococcal tonsillopharyngitis, 97

Peptidyl-dipeptidase A

Losartan did not improve survival more than did captopril but was better tolerated in symptomatic heart failure, 87

Pharyngitis

Review: Twice-daily dosing of penicillin V is as effective as more frequent dosing for streptococcal tonsillopharyngitis, 97

Phlebotomy

Pacifiers have greater analgesic effect than sweet solutions during venipuncture in full-term newborns, 25

Physical examination

Clinical assessment of the reliability of the examination (CARE) (Editorial), A-11-12 (No. 2)

Physical fitness

Review: Brief primary care interventions are moderately effective for increasing physical activity, 104

Physical therapy

Manual physical therapy and exercise improved function in osteoarthritis of the knee, 57

Physician-patient relations

Review: Providing recordings or summaries of consultations may help patients with cancer, 28

Piperidines

Sibrafiban was more toxic than aspirin for prevention of cardiovascular events after acute coronary syndromes, 5

Platelet aggregation inhibitors

Eptifibatide was cost-effective in patients with unstable angina or non-Q-wave myocardial infarction, 76

Review: High- and moderate-dose oral anticoagulants reduce events in CAD but increase major bleeding and are no more effective than aspirin, 8

Platelet glycoprotein GPIIb-IIIa complex

Eptifibatide was cost-effective in patients with unstable angina or non-Q-wave myocardial infarction, 76

Sibrafiban was more toxic than aspirin for prevention of cardiovascular events after acute coronary syndromes, 5

Pneumonia

A critical pathway reduced resource use without compromising safety and effectiveness in community-acquired pneumonia, 72

Invasive diagnosis for ventilator-associated pneumonia reduced 14-day mortality and antibiotic use, 93

Tachypnea is a useful predictor of pneumonia in children with acute respiratory infection, 68

Postmenopausal osteoporosis

Risedronate was effective and well tolerated in postmenopausal women with osteoporosis, 15

Postoperative complications

Routine testing before cataract surgery did not reduce medical adverse events, 60

Practice guideline

Practice-based education did not increase recognition of depression by primary-care physicians nor improve the outcome of depression, 74

Prednisone

Inhaled budesonide and prednisone led to similar rates of asthma relapse after emergency department discharge, 51

Pregnancy complications, cardiovascular

Review: Treatment-induced blood pressure reductions in pregnancy may be associated with decreased fetal growth, 42

Preoperative care

Routine testing before cataract surgery did not reduce medical adverse events, 60

Primary health care

Practice-based education did not increase recognition of depression by primary-care physicians nor improve the outcome of depression, 74

Review: Brief primary care interventions are moderately effective for increasing physical activity, 104

Propafenone

Low-dose amiodarone was better than sotalol or propafenone for preventing first recurrence of atrial fibrillation, 88

Prostatectomy

Pelvic floor re-education reduced incontinence at 1 year after radical prostatectomy, 108

Prostatic neoplasms

Antiandrogen therapy immediately after surgery for prostate cancer improved survival, 29

Pulmonary embolism

Hormone replacement therapy was associated with increased venous thromboembolism and deep venous thrombosis, 106

- Low-molecular-weight heparin reduced recurrent VTE in patients with pulmonary embolism and proximal DVT, 6
- Review: Helical computed tomography for suspected pulmonary embolism needs further evaluation, 32-33
- Review: Low-molecular-weight heparin reduces death but not recurrent VTE events, bleeding, or thrombocytopenia in VTE, 7
- SimpliRED D-dimer assay did not rule out venous thromboembolism in the emergency department, 66
- Pyrimethamine
- Adding 3-dose artesunate to pyrimethamine-sulfadoxine reduced treatment failure in children with acute uncomplicated malaria, 98
- ## Q
- Quality indicators
- Review: High-volume hospitals had lower hospital mortality rates than did low-volume hospitals for various procedures, 119
- Quinolones
- Oral montelukast was better than inhaled salmeterol for reducing exercise-induced bronchoconstriction in adults with asthma, 3
- ## R
- Reagent strips
- The Uriscreeen test was not better than standard urinalysis and dipstick tests for detecting urinary tract infection in children, 35
- Recombinant proteins
- Intra-arterial recombinant prourokinase improved neurologic recovery in acute middle cerebral artery ischemic stroke, 19
- Referral and consultation
- Review: High-volume hospitals had lower hospital mortality rates than did low-volume hospitals for various procedures, 119
- Religion
- Review: "Distant healing" is often effective for improving patient outcomes, 107
- Renal artery obstruction
- Angioplasty was no better than medication for reducing blood pressure or slowing disease progression in renal artery stenosis, 92
- Respiration disorders
- A respiratory rate of  $\geq 60$  breaths per minute had high sensitivity for detecting hypoxia in infants, 69
- Respiration, artificial
- Review: Quality of sedation in the ICU is similar for propofol and midazolam, but propofol shortens the time to extubation, 50
- Reverse transcriptase inhibitors
- Efavirenz plus zidovudine and lamivudine was more effective than indinavir plus zidovudine and lamivudine in adults with HIV-1 infection, 22
- Rheumatic diseases
- Review: Silicone breast implants do not increase the risk for connective-tissue diseases, 118
- Riluzole
- Review: Riluzole increases 12-month survival in amyotrophic lateral sclerosis, 54
- ## S
- Sarcoidosis, pulmonary
- Review: Oral but not inhaled corticosteroids improve radiographic outcomes in patients with pulmonary sarcoidosis, 102
- Sex factors
- Women with unstable angina received fewer cardiac procedures than did men but were less likely to develop other cardiac events, 70
- Signs and symptoms
- Clinical assessment of the reliability of the examination (CARE) (Editorial), A-11-12 (No. 2)
- Sleep disorders
- Review: Long-term effectiveness of nondrug treatments for sleep problems in young children is more evident, 61
- Social support
- Methadone maintenance was more effective for treatment retention of opioid dependence than was psychosocially enriched detoxification, 109
- Sotalol
- Low-dose amiodarone was better than sotalol or propafenone for preventing first recurrence of atrial fibrillation, 88
- Spinal cord diseases
- Review: Antimicrobial prophylaxis reduces asymptomatic bacteriuria in patients with neurogenic bladder, 21
- Spouses
- Older adults who reported strain when caring for a spouse with disabilities had increased mortality, 37
- Stress, psychological
- Older adults who reported strain when caring for a spouse with disabilities had increased mortality, 37
- Surgical procedures, operative
- Review: High-volume hospitals had lower hospital mortality rates than did low-volume hospitals for various procedures, 119
- Syncope
- A new arrival: evidence about differential diagnosis (Editorial), A-11-12 (No. 3)
- A simple algorithm improved diagnostic performance for patients presenting with syncope, 121
- Syndrome
- Review: Antidepressants are effective for clinical improvement in unexplained physical symptoms and syndromes, 53
- ## T
- Tachycardia, ventricular
- The implantable cardioverter defibrillator did not reduce mortality in ventricular arrhythmia, 43
- Tape recording
- Review: Providing recordings or summaries of consultations may help patients with cancer, 28
- Telephone
- Feedback to physicians plus telephone care management improved outcomes in primary care patients with depression, 73
- Tennis elbow
- Local injection, naproxen, and simple analgesia led to similar 1-year rates of symptom relief in lateral epicondylitis, 26
- Therapeutic touch
- Review: "Distant healing" is often effective for improving patient outcomes, 107
- Thiazoles
- Low-dose rosiglitazone plus sulfonylurea was effective and safe for treating hyperglycemia in type 2 diabetes mellitus, 49

- Metformin plus rosiglitazone had more effective glycemic control than metformin alone in type 2 diabetes mellitus, 111
- Thiophenes  
Review: Thienopyridines are more effective than aspirin for preventing vascular events in high-risk patients, 45
- Thromboembolism  
The recurrence rate of venous thromboembolism after a first or second episode of deep venous thrombosis was high, 117
- Thrombolytic therapy  
Intra-arterial recombinant prourokinase improved neurologic recovery in acute middle cerebral artery ischemic stroke, 19  
Recombinant tissue-type plasminogen activator at 3 to 5 hours after ischemic acute stroke onset was not effective or safe, 18
- Ticlopidine  
Review: Thienopyridines are more effective than aspirin for preventing vascular events in high-risk patients, 45
- Tiotropium  
Tiotropium improved lung function more than did ipratropium in chronic obstructive pulmonary disease, 101
- Tissue plasminogen activator  
Recombinant tissue-type plasminogen activator at 3 to 5 hours after ischemic acute stroke onset was not effective or safe, 18
- Tomography, x-ray computed  
CT scanning before fiberoptic bronchoscopy improved the accuracy of bronchoscopy in suspected endobronchial carcinoma, 100  
Review: Helical computed tomography for suspected pulmonary embolism needs further evaluation, 32-33
- Tonsillitis  
Review: Twice-daily dosing of penicillin V is as effective as more frequent dosing for streptococcal tonsillopharyngitis, 97
- Trichomonas infections  
Review: Papanicolaou and wet-mount smears have low sensitivity but high specificity for detecting vaginal trichomoniasis, 113
- U**
- Upper respiratory tract infections  
Review: Diagnostic tests for bacterial rhinosinusitis have moderate accuracy; antibiotics reduce clinical failures, 31
- Urinalysis  
The Uriscreeen test was not better than standard urinalysis and dipstick tests for detecting urinary tract infection in children, 35
- Urinary incontinence  
Pelvic floor re-education reduced incontinence at 1 year after radical prostatectomy, 108
- Urinary plasminogen activator  
Intra-arterial recombinant prourokinase improved neurologic recovery in acute middle cerebral artery ischemic stroke, 19

- Urinary tract infections  
Review: Antimicrobial prophylaxis reduces asymptomatic bacteriuria in patients with neurogenic bladder, 21  
The Uriscreeen test was not better than standard urinalysis and dipstick tests for detecting urinary tract infection in children, 35

**V**

- Vaginal smears  
Review: The extended-tip spatula is more effective than the Ayre's spatula for detecting karyosis in cervical smears, 30  
Review: Newer technologies improve sensitivity in detecting cervical lesions but at a substantially increased cost, 34  
Review: Papanicolaou and wet-mount smears have low sensitivity but high specificity for detecting vaginal trichomoniasis, 113
- Venous thrombosis  
Hormone replacement therapy was associated with increased venous thromboembolism and deep venous thrombosis, 106  
Low-molecular-weight heparin reduced recurrent VTE in patients with pulmonary embolism and proximal DVT, 6  
The recurrence rate of venous thromboembolism after a first or second episode of deep venous thrombosis was high, 117  
Review: Helical computed tomography for suspected pulmonary embolism needs further evaluation, 32-33  
Review: Low-molecular-weight heparin reduces death but not recurrent VTE events, bleeding, or thrombocytopenia in VTE, 7  
SimpliRED D-dimer assay did not rule out venous thromboembolism in the emergency department, 66
- Ventilators, mechanical  
Invasive diagnosis for ventilator-associated pneumonia reduced 14-day mortality and antibiotic use, 93
- Ventricular dysfunction, left  
A combination of tests done in general practice could assess the risk for left ventricular systolic dysfunction, 114
- Verapamil  
Verapamil reduced attack frequency and the use of abortive agents in episodic cluster headache, 105
- Vitamin E  
Vitamin E did not reduce myocardial infarction, death, or stroke in adults at high risk for cardiovascular events, 47

**W**

- Weight loss  
Weight loss in obese patients with asthma improved lung function and health status, 103

